

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

^PrZYDELIG[®]

idelalisib tablets

Tablet, 150 mg and 100 mg, oral

Antineoplastic Agent

Gilead Sciences Canada, Inc.
Mississauga, ON L5N 2W3

www.gilead.ca

Submission Control No.: 240413

Date of Initial Approval: 19 February
2015

Date of Revision: 18 August 2020

Table of Contents

PART I. HEALTH PROFESSIONAL INFORMATION 3

 SUMMARY PRODUCT INFORMATION..... 3

 INDICATIONS AND CLINICAL USE 3

 CONTRAINDICATIONS 4

 WARNINGS AND PRECAUTIONS 5

 ADVERSE REACTIONS..... 10

 DRUG INTERACTIONS..... 17

 DOSAGE AND ADMINISTRATION 18

 OVERDOSAGE..... 21

 ACTION AND CLINICAL PHARMACOLOGY 21

 STORAGE AND STABILITY 24

 SPECIAL HANDLING INSTRUCTIONS..... 24

 DOSAGE FORMS, COMPOSITION AND PACKAGING 24

PART II. SCIENTIFIC INFORMATION 25

 PHARMACEUTICAL INFORMATION..... 25

 CLINICAL TRIALS 26

 DETAILED PHARMACOLOGY 31

 TOXICOLOGY 31

 REFERENCES..... 34

PATIENT MEDICATION INFORMATION 35

PrZYDELIG®

(idelalisib)

PART I. HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form/Strength	Clinically Relevant Nonmedicinal Ingredients
Oral	Tablet Idelalisib 150 mg Idelalisib 100 mg	microcrystalline cellulose, hydroxypropyl cellulose, croscarmellose sodium, sodium starch glycolate, and magnesium stearate. <i>For a complete listing, see the DOSAGE FORMS, COMPOSITION, AND PACKAGING section.</i>

INDICATIONS AND CLINICAL USE

Chronic Lymphocytic Leukemia

ZYDELIG (idelalisib) is indicated in combination with rituximab for the treatment of patients with relapsed chronic lymphocytic leukemia (CLL).

Effectiveness of ZYDELIG in combination with rituximab is based on progression free survival benefit with limited follow up in a study of patients who were not fit to receive cytotoxic therapy.

Refer to the Rituxan Product Monograph for rituximab product information.

Follicular Lymphoma

ZYDELIG (idelalisib) is indicated as a monotherapy for the treatment of patients with follicular lymphoma who have received at least two prior systemic regimens and are refractory to both rituximab and an alkylating agent.

Geriatrics (≥65 years of age):

In clinical studies of ZYDELIG in patients with follicular lymphoma or CLL, no major differences in effectiveness were observed in patients 65 years of age or older compared with younger patients. Adverse events were more common and led more commonly to

negative outcomes in patients over the age of 65 years (see **WARNINGS and PRECAUTIONS**).

Pediatrics (<18 years of age):

Safety and effectiveness in children less than 18 years of age have not been established.

CONTRAINDICATIONS

ZYDELIG is contraindicated in first-line CLL and early-line indolent non-Hodgkin lymphoma (iNHL) outside a clinical trial. See **WARNINGS AND PRECAUTIONS, Serious Infections** section below.

ZYDELIG is contraindicated in patients with known hypersensitivity to any of the components of the product. For a complete listing, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING** section of the Product Monograph.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

ZYDELIG® should only be prescribed by a qualified physician who is experienced in the use of anti-cancer agents.

Prophylaxis for *Pneumocystis carinii/jirovecii* pneumonia (PCP/PJP) and monitoring for cytomegalovirus (CMV) are required during treatment with ZYDELIG.

The following are clinically significant adverse events:

- Serious infections, including fatal cases (see **Serious Infections** below)
- Hepatotoxicity (see **Hepatic** below)
- Severe diarrhea/colitis, including fatal cases (see **Gastrointestinal** below)
- Pneumonitis, including fatal cases (see **Respiratory** below)
- Severe cutaneous reactions, including drug reaction with eosinophilia and systemic symptoms (DRESS), and fatal cases of Stevens-Johnson Syndrome and Toxic Epidermal Necrolysis (see **Skin** below)

Carcinogenesis and Mutagenesis

The carcinogenicity potential of idelalisib was evaluated in a 26-week transgenic RasH2 mouse study and a 2-year rat study. A small increase in pancreatic islet cell tumors was noted in male rats (see **TOXICOLOGY, Carcinogenesis**). ZYDELIG demonstrated no significant mutagenicity or genotoxicity *in vitro*, but demonstrated genotoxicity *in vivo* at a high dose (see **TOXICOLOGY, Mutagenesis**).

Drug Interactions

Cytochrome P450 (CYP) interactions

ZYDELIG is a strong CYP3A inhibitor. Coadministration of ZYDELIG with CYP3A substrates may increase their systemic exposures. Caution is recommended if ZYDELIG is coadministered with narrow therapeutic index CYP3A substrates (see **DRUG INTERACTIONS**).

Gastrointestinal

Diarrhea/Colitis:

Cases of severe diarrhea/colitis were reported commonly and occurred relatively late (months) after the start of therapy with ZYDELIG. Severe diarrhea due to ZYDELIG responds poorly to antimotility agents. Most cases resolved within a few weeks with drug interruption and additional symptomatic treatment (e.g., anti-inflammatory corticosteroid

agents such as enteric budesonide) but some had a fatal outcome (see **DOSAGE AND ADMINISTRATION, Dose Modification**). Severe diarrhea/colitis occurring with administration of ZYDELIG has been associated with dehydration which has responded to intravenous fluid and electrolyte replacement. Causes of diarrhea related to gastrointestinal infection should be ruled out.

For CTCAE Grade 1 diarrhea/colitis, provide anti-diarrheal (e.g., loperamide) and maintain ZYDELIG dose. For Grade 2 diarrhea/colitis, withhold ZYDELIG and monitor at least weekly until resolved to Grade ≤ 1 .

For CTCAE Grade 3 or 4 diarrhea/colitis, withhold ZYDELIG. Consider addition of anti-inflammatory agent (e.g., sulfasalazine, budesonide). Monitor at least weekly until resolved to Grade ≤ 1 , then may resume ZYDELIG at 100 mg BID.

Treatment of patients with ongoing inflammatory bowel disease is not recommended.

Hematologic

Neutropenia:

Treatment-emergent Grade 3 or 4 neutropenia and cases of febrile neutropenia, some which have been fatal, have occurred in patients treated with ZYDELIG. Monitor blood counts in all patients at least every 2 weeks for the first 6 months of therapy with ZYDELIG, and at least weekly in patients while absolute neutrophil counts are less than $1.0 \times 10^9/L$ (see **DOSAGE AND ADMINISTRATION, Dose Modification**).

Hepatic

Hepatotoxicity:

Elevations in ALT and AST Grade 3 or 4 (greater than 5 times the upper limit of normal) have been observed in clinical trials of ZYDELIG. These laboratory findings were generally observed within the first 12 weeks of treatment, asymptomatic, and reversible within 3-4 weeks with dose interruption. While most patients resumed treatment at a lower dose, recurrence of ALT and AST elevations were common (see **DOSAGE AND ADMINISTRATION, Dose Modification**). Monitor ALT, AST, and total bilirubin in all patients every 2 weeks for the first 3 months of treatment, then every 1 to 3 months thereafter, and as clinically indicated.

For CTCAE Grade 1 (ALT/AST $\leq 3 \times$ ULN) or Grade 2 (ALT/AST $> 3-5 \times$ ULN), maintain ZYDELIG dose. Monitor at least weekly until ALT/AST are $\leq 1 \times$ ULN.

For CTCAE Grade 3 (ALT/AST >5-20 x ULN) or Grade 4 (ALT/AST >20 x ULN), withhold ZYDELIG. Monitor at least weekly until ALT/AST are ≤ 1 x ULN, then may resume ZYDELIG at 100 mg BID.

Discontinue ZYDELIG for recurrent hepatotoxicity.

Treatment of patients with active hepatitis or liver disease is not recommended.

Immune

Anaphylaxis:

Serious allergic reactions, including anaphylaxis, have been reported in patients on ZYDELIG. In patients who develop serious allergic reactions, discontinue ZYDELIG permanently and institute appropriate supportive measures.

Serious Infections:

Treatment with ZYDELIG should not be initiated in patients with any evidence of ongoing systemic bacterial, fungal or viral infection.

Serious and fatal infections have occurred with ZYDELIG, including opportunistic infections such as *Pneumocystis carinii/jirovecii* pneumonia (PCP/PJP) and cytomegalovirus (CMV).

An increase in serious adverse events and deaths, primarily due to infections, was observed in patients receiving ZYDELIG compared with the control arms in a first-line study of CLL and two studies of relapsed early-line iNHL. Based on these findings, and the unfavourable benefit/risk assessment at interim analysis, these studies and all first-line studies in patients with CLL or iNHL were terminated. ZYDELIG is contraindicated in first-line CLL and early-line iNHL outside a clinical trial.

Administer prophylaxis for PCP/PJP to all patients throughout ZYDELIG treatment and for a period of 2 to 6 months after discontinuation. The duration of post-treatment prophylaxis should be based on clinical judgment and may take into account a patient's risk factors such as concomitant corticosteroid treatment and prolonged neutropenia.

Regular clinical and laboratory monitoring for CMV infection should be conducted. Treatment with ZYDELIG should be permanently discontinued if there is evidence of CMV infection or viremia (positive polymerase chain reaction (PCR) or antigen test).

Cases of progressive multifocal leukoencephalopathy (PML) have been reported following the use of ZYDELIG within the context of prior- or concomitant immunosuppressive therapies that have been associated with PML. Physicians should consider PML in the differential diagnosis in patients with new or worsening neurological, cognitive or

behavioral signs or symptoms. If PML is suspected then appropriate diagnostic evaluations should be undertaken and treatment suspended until PML is excluded. If any doubt exists, referral to a neurologist and appropriate diagnostic measures for PML including MRI scan preferably with contrast, cerebrospinal fluid (CSF) testing for JC viral DNA and repeat neurological assessments should be considered. Discontinue ZYDELIG permanently in patients with confirmed PML.

Neurologic

Effects on Ability to Drive and Use Machines

No studies of the effects of ZYDELIG on the ability to drive or use machines have been performed. A detrimental effect on such activities is not expected based on the known pharmacology and safety profile of ZYDELIG.

Respiratory

Pneumonitis:

Cases of pneumonitis, including organizing pneumonia, some with fatal outcome, have occurred with ZYDELIG. Time to occurrence of pneumonitis after the start of therapy with ZYDELIG was highly variable, ranging from a few weeks to over one year. Patients should be monitored for respiratory signs and symptoms throughout treatment and should be advised to report new respiratory symptoms promptly. In patients presenting with serious lung adverse events, ZYDELIG should be interrupted and the patient assessed for an explanatory etiology. In patients with drug-related pneumonitis or organizing pneumonia, ZYDELIG should be permanently discontinued and appropriate treatment with systemic corticosteroids should be initiated (see **DOSAGE AND ADMINISTRATION, Dose Modification**).

Skin

Severe Cutaneous Reactions:

Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug reaction with eosinophilia and systemic symptoms (DRESS) have occurred with ZYDELIG. Cases of SJS and TEN with fatal outcomes have been reported when ZYDELIG was administered concomitantly with other medications associated with these syndromes. If SJS, TEN, or DRESS is suspected, ZYDELIG should be interrupted and the patient treated accordingly. If SJS, TEN, or DRESS is confirmed, permanently discontinue ZYDELIG.

Other severe or life-threatening (Grade ≥ 3) cutaneous reactions, including dermatitis exfoliative, rash, rash erythematous, rash generalized, rash macular, rash maculo-papular, rash papular, rash pruritic, exfoliative rash, and skin disorder, have been reported in

ZYDELIG-treated patients. If patients develop severe cutaneous reactions, discontinue ZYDELIG (see **DOSAGE AND ADMINISTRATION, Dose Modification**).

For CTCAE Grade 1 rash, maintain ZYDELIG dose. For Grade 2 rash, withhold ZYDELIG until Grade ≤ 1 .

For CTCAE Grade 3 or 4 rash, withhold ZYDELIG. Monitor at least weekly until resolved to Grade ≤ 1 , then may resume ZYDELIG at 100 mg BID.

Photosensitivity:

In vitro studies have shown that idelalisib has phototoxic potential (see **TOXICOLOGY**). Patients should be advised to avoid sun exposure or wear sufficient sun protection.

Special Populations

Pregnant Women:

There are no adequate and well-controlled studies of ZYDELIG in pregnant women.

Based on findings in animals (see **TOXICOLOGY**), idelalisib may cause fetal harm when administered to a pregnant woman. In studies of pregnant rats, idelalisib treatment was associated with increased post-implantation loss, decreased fetal weights, and skeletal malformations.

ZYDELIG should not be used during pregnancy. If this drug is used during pregnancy, or if the patient becomes pregnant while receiving this drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing potential should be advised to avoid becoming pregnant while receiving ZYDELIG. Women of child bearing potential should use highly effective contraceptive measures while taking ZYDELIG and for 1 month after stopping treatment. Idelalisib may reduce the effectiveness of hormonal contraceptives (see **DRUG INTERACTIONS**). Women who use hormonal methods of birth control should add a barrier method.

Nursing Women:

It is not known whether idelalisib or its metabolites are excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from ZYDELIG, discontinue nursing when taking ZYDELIG.

Hepatic Impairment:

Intensified monitoring of adverse events is recommended in patients with impaired hepatic function as exposure is expected to be increased in this population. The AUC of idelalisib increased up to 1.7-fold in subjects with ALT or AST or bilirubin greater than the upper limit

of normal (ULN) compared to healthy subjects with normal ALT or AST or bilirubin values. Safety and efficacy data are not available in patients with baseline ALT or AST values greater than 2.5 x ULN or bilirubin values greater than 1.5 x ULN, as these patients were excluded from pivotal trials.

Geriatrics:

The incidence of \geq Grade 3 AEs was higher among subjects \geq 65 years of age compared with subjects < 65 years of age (79.0% vs 63.2%). Subjects \geq 65 years of age had a higher incidence of idelalisib discontinuation due to an AE compared with subjects < 65 years of age (24.6% vs 15.7%). A higher incidence of SAEs was observed in subjects \geq 65 years of age compared with subjects < 65 years of age (59.9% vs 40.0% for all subjects).

Monitoring and Laboratory Tests

Hepatic function tests (ALT, AST, and total bilirubin) should be measured every 2 weeks for the first 3 months of treatment, then every 1 to 3 months thereafter, and as clinically indicated (see **Hepatic** section above).

Monitor blood counts at least every 2 weeks for the first 6 months of therapy, and at least weekly in patients while absolute neutrophil counts are less than $1.0 \times 10^9/L$. Monitor all patients for CMV infection during treatment with ZYDELIG.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

ZYDELIG is associated with infections. Higher frequencies of infections overall, including Grade 3 and 4 infections, were observed in the ZYDELIG arms compared to the control arms of ZYDELIG clinical studies. Most frequently observed were infections in the respiratory system and septic events. In many instances the pathogen was not identified; however, both conventional and opportunistic pathogens, including PCP/PJP and CMV, were among those identified. Nearly all PCP/PJP infections, including fatal cases, occurred in the absence of PCP/PJP prophylaxis. There have been cases of PCP/PJP after stopping idelalisib treatment.

Chronic Lymphocytic Leukemia

In the Phase 3 study in CLL, 220 previously treated patients were randomised to receive ZYDELIG (150 mg BID) + rituximab or placebo + rituximab. Serious adverse reactions were reported in 54 (49%) patients treated with ZYDELIG + rituximab. The most frequent (\geq 2%) serious adverse reactions reported for patients treated with ZYDELIG were pneumonia (17%), pyrexia (9%), sepsis (8%), febrile neutropenia (5%), and diarrhea (5%). Adverse

reactions that led to discontinuation of ZYDELIG occurred in 11 (10%) patients. The most common adverse reactions that led to treatment discontinuations were hepatotoxicity and diarrhea/colitis. A total of 39 patients (35%) had dose interruptions, 16 patients (15%) had dose reductions, and 11 patients (10%) had drug discontinuation due to adverse reactions. Patients may have had more than 1 type of dose modification. The most common reasons for dose reductions were elevated transaminases, diarrhea, and neutropenia.

Indolent Non-Hodgkin Lymphoma

In the Phase 1 and 2 studies in previously treated iNHL, 146 patients were treated with ZYDELIG monotherapy at a dose of 150 mg BID. Serious adverse reactions were reported in 73 (50%) patients treated with ZYDELIG. The most frequent serious adverse reactions were pneumonia (15%), diarrhea (11%), and pyrexia (9%). Among the 146 iNHL patients who received ZYDELIG 150 mg BID as a single agent, sixty-two (43%) had dose interruptions, thirty-four (23%) had dose reductions, and thirty-six (25%) had drug discontinuation due to adverse reactions. Patients may have had more than 1 type of dose modification. The most common reasons for dose modifications were diarrhea, elevated transaminases, and neutropenia.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Table 1 summarizes adverse events reported for ZYDELIG + rituximab and placebo + rituximab arms in 218 patients with CLL. Adverse events reflect exposure to ZYDELIG with a median duration of 5.0 months and exposure to placebo with a median duration of 3.7 months. Adverse events occurring at 5% incidence and 2% greater in the ZYDELIG arm when compared to the placebo arm are provided in Table 1.

Table 1 Adverse Events Reported in ≥5% of Patients with CLL and Occurred at ≥2% Higher Incidence in Patients Treated with ZYDELIG

Adverse Event	ZYDELIG + R N=110 (%)		Placebo + R N=108 (%)	
	Any Grade	Grade ≥3	Any Grade	Grade ≥3
Gastrointestinal disorders				

Adverse Event	ZYDELIG + R N=110 (%)		Placebo + R N=108 (%)	
	Any Grade	Grade ≥3	Any Grade	Grade ≥3
nausea	28 (25)	0	23 (21)	0
vomiting	14 (13)	0	9 (8)	0
diarrhea ^(a)	23 (21)	6 (5)	17 (16)	0
gastroesophageal reflux disease	7 (6)	0	1 (1)	0
stomatitis	7 (6)	2 (2)	2 (2)	0
Nervous system disorders				
headache	11 (10)	1 (1)	5 (5)	0
General disorders and administration site conditions				
pyrexia	38 (35)	3 (3)	18 (17)	1 (1)
chills	23 (21)	2 (2)	17 (16)	0
pain	8 (7)	0	2 (2)	0
Skin and subcutaneous tissue disorders				
rash ^(b)	20 (18)	4 (4)	7 (6)	1 (1)
Respiratory, thoracic, and mediastinal disorders				
pneumonia ^(c)	25 (23)	18 (16)	19 (18)	14 (13)
nasal congestion	6 (5)	0	2 (2)	0
Infections and infestations				
sepsis ^(d)	9 (8)	8 (7)	4 (4)	4 (4)
bronchitis	7 (6)	1 (1)	3 (3)	1 (1)
sinusitis	9 (8)	0	4 (4)	0
urinary tract infection	6 (5)	0	3 (3)	2 (2)
Musculoskeletal and connective tissue disorders				
arthralgia	8 (7)	1 (1)	4 (4)	1 (1)

(a) Diarrhea includes the following preferred terms: diarrhea, colitis.

(b) Rash includes the following preferred terms: dermatitis exfoliative, rash, rash macular, rash maculo-papular, rash papular, rash pruritic, and skin disorder.

(c) Pneumonia includes the terms: pneumonia, pneumonitis, lung infection, lung infiltration, pneumocystis jiroveci pneumonia, pneumonia legionella, lung infection pseudomonal, pneumonia fungal, respiratory tract infection, lower respiratory tract infection, and lower respiratory tract infection bacterial.

(d) Sepsis includes the terms: sepsis, septic shock, neutropenic sepsis, and sepsis syndrome.

R: rituximab

Table 2 summarizes adverse events in 146 patients with iNHL treated with ZYDELIG 150 mg BID. ZYDELIG is indicated for use in patients with follicular lymphoma. Adverse events reflect exposure to ZYDELIG with a median duration of 6.1 months.

Table 2 Adverse Events Reported in ≥ 10% Patients with Indolent Non-Hodgkin Lymphoma Treated with ZYDELIG 150 mg BID

Adverse Event	ZYDELIG Monotherapy N=146 (%)	
	Any Grade	Grade ≥3
Gastrointestinal disorders		
diarrhea ^(a)	68 (47)	20 (14)
nausea	42 (29)	2 (1)
abdominal pain ^(b)	38 (26)	3 (2)
vomiting	22 (15)	2 (1)
General disorders and administration site conditions		
fatigue	44 (30)	2 (1)
pyrexia	41 (28)	3 (2)
asthenia	17 (12)	3 (2)
peripheral edema	15 (10)	3 (2)
Infections and infestations		
upper respiratory tract infection	18 (12)	0
Respiratory, thoracic, and mediastinal disorders		
pneumonia ^(c)	37 (25)	23 (16)
cough	42 (29)	1 (1)
dyspnea	25 (17)	6 (4)
Skin and subcutaneous disorders		
rash ^(d)	31 (21)	4 (3)
night sweats	18 (12)	0
Nervous system disorders		
headache	16 (11)	1 (1)
Metabolism and nutrition disorders		
decreased appetite	24 (16)	1 (1)
Psychiatric disorders		
insomnia	17 (12)	0

(a) Diarrhea includes the following preferred terms: diarrhea, colitis, enterocolitis, and gastrointestinal inflammation.

(b) Abdominal pain includes the following preferred terms: abdominal pain, abdominal pain upper, abdominal pain lower, and abdominal discomfort.

(c) Pneumonia includes the terms: pneumonia, pneumonitis, interstitial lung disease, lung infiltration, pneumonia aspiration, respiratory tract infection, atypical pneumonia, lung infection, pneumocystis jiroveci pneumonia, bronchopneumonia, pneumonia necrotizing,

lower respiratory tract infection, pneumonia pneumococcal, pneumonia staphylococcal, pneumonia streptococcal, pneumonia cytomegaloviral, and respiratory syncytial virus infection.

(d) Rash includes the following preferred terms: dermatitis exfoliative, rash, rash erythematous, rash macular, rash maculo-papular, rash pruritic, and exfoliative rash.

Abnormal Hematologic and Clinical Chemistry Findings

Laboratory Abnormalities

Table 3 summarizes the treatment-emergent laboratory abnormalities reported for ZYDELIG + rituximab and placebo + rituximab arms in patients with CLL.

Table 3 Treatment-emergent Laboratory Abnormalities Reported in ≥10% of Patients with CLL Occurring at a ≥5% Higher Incidence in Patients Receiving ZYDELIG

Laboratory Parameter	ZYDELIG + R N=110 (%)		Placebo + R N=108 (%)	
	Any Grade	Grade 3–4	Any Grade	Grade 3–4
Hematology abnormalities				
neutrophil count decreased	66 (60)	41 (37)	55 (51)	29 (27)
lymphocyte count decreased	22 (20)	10 (9)	13 (12)	4 (4)
lymphocyte count increased	27 (25)	20 (18)	10 (9)	5 (5)
Serum chemistry abnormalities				
ALT increased	38 (35)	9 (8)	11 (10)	1 (1)
AST increased	27 (25)	6 (5)	15 (14)	0
GGT increased	29 (26)	2 (2)	15 (14)	3 (3)
triglycerides (hypertriglyceridemia)	62 (56)	3 (3)	37 (34)	0
hyperglycemia	59 (54)	8 (7)	50 (46)	2 (2)
hypoglycemia	12 (11)	0	5 (5)	0
hyponatremia	22 (20)	2 (2)	16 (15)	7 (6)

Grades were obtained per CTCAE version 4.03.

R: rituximab

Table 4 summarizes the treatment-emergent laboratory abnormalities in patients with iNHL treated with ZYDELIG 150 mg BID.

Table 4 Treatment-emergent Laboratory Abnormalities in Patients with Indolent Non-Hodgkin Lymphoma Treated with ZYDELIG 150 mg BID

Laboratory Abnormality	ZYDELIG Monotherapy N=146 (%)	
	Any Grade	Grade 3-4
Serum chemistry abnormalities		
ALT increased	73 (50)	27 (18)
AST increased	60 (41)	18 (12)
neutrophils decreased	78 (53)	36 (25)
hemoglobin decreased	41 (28)	3 (2)
platelets decreased	38 (26)	9 (6)

Grades were obtained per CTCAE version 4.03.

Post-Market Adverse Drug Reactions

In addition to adverse reactions from clinical studies, the following adverse reactions have been identified during post-approval use of ZYDELIG. Because these reactions were reported voluntarily from a population of unknown size, estimates of frequency cannot be made.

Skin and subcutaneous tissue disorders:

Stevens-Johnson Syndrome (SJS), Toxic Epidermal Necrolysis (TEN), and drug reaction with eosinophilia and systemic symptoms (DRESS) (see **WARNINGS and PRECAUTIONS, Skin**)

Respiratory, Thoracic and Mediastinal disorders:

Organizing pneumonia (see **WARNINGS and PRECAUTIONS, Respiratory**)

Infections and infestations:

Progressive multifocal leukoencephalopathy (see **WARNINGS and PRECAUTIONS, Serious infections**)

DRUG INTERACTIONS

Drug-Drug Interactions

Idelalisib is metabolized primarily via aldehyde oxidase, and to a lesser extent via CYP3A and glucuronidation (UGT1A4). The primary circulating metabolite (GS-563117) exceeds idelalisib human plasma levels at steady-state. GS-563117 is inactive against PI3K δ .

Effects of Other Drugs on ZYDELIG

CYP3A Inducers

A clinical drug interaction study found that coadministration of ZYDELIG with rifampin (a strong CYP3A inducer) resulted in a ~75% reduction in idelalisib plasma AUC_{inf}. Coadministration of ZYDELIG with strong CYP3A inducers such as rifampin, phenytoin, St. John's Wort, or carbamazepine should be avoided.

CYP3A Inhibitors

A clinical drug interaction study found that coadministration of ZYDELIG with ketoconazole (a strong CYP3A inhibitor) resulted in a 26% increase in ZYDELIG C_{max} and a 79% increase in AUC_{inf}, indicating that ZYDELIG is not a sensitive CYP3A substrate. If patients are taking concomitant strong CYP3A inhibitors, monitor for signs of toxicity.

Other

Co-administration of drugs which are aldehyde oxidase inhibitors (e.g. raloxifene) could increase idelalisib plasma concentrations. A clinical drug interaction study has not been conducted.

Effects of ZYDELIG on Other Drugs

CYP3A Substrates

In vitro, the major circulating metabolite was shown to be an irreversible inhibitor of CYP3A. Return to normal CYP3A enzyme activity is therefore expected to take several days after stopping idelalisib administration. A clinical drug interaction study found that coadministration of ZYDELIG with midazolam (a sensitive CYP3A substrate) resulted in a ~140% increase in C_{max} and a ~440% increase in AUC_{inf} of midazolam. Accordingly, ZYDELIG is considered to be a strong CYP3A inhibitor. Coadministration of ZYDELIG with CYP3A substrates (e.g., certain antiarrhythmics, calcium channel blockers, benzodiazepines, HMG-CoA reductase inhibitors, phosphodiesterase-5 (PDE5) inhibitors, and warfarin) may increase their systemic exposures.

Caution is recommended if ZYDELIG is coadministered with narrow therapeutic index CYP3A substrates (e.g., alfentanil, cyclosporine, sirolimus, tacrolimus, pimozide, fentanyl, quinidine, ergotamine, dihydroergotamine).

CYP2C8 Substrates

In vitro, idelalisib inhibited CYP2C8. *In vivo* studies to investigate the clinical relevance have not been conducted. Caution is advised when co-administering idelalisib with narrow therapeutic index CYP2C8 substrates (e.g. repaglinide).

Substrates of CYP2B6, CYP2C9 and CYP2C19

In vitro, idelalisib demonstrated the potential to induce CYP2B6, CYP2C8 and CYP2C9, and based on these findings, is likely to induce CYP2C19. Clinical drug interaction studies have not been conducted. Caution is advised upon co-administration of idelalisib with substrates of these enzymes with narrow therapeutic indices (warfarin, phenytoin, S-mephenytoin).

Hormonal Contraceptives

In vitro, idelalisib induces CYP3A4, CYP2C9 and UGT1A1. All of these enzymes are involved in the first-pass metabolism of ethinyl estradiol in the gut wall and liver. A clinical drug interaction study has not been conducted, however idelalisib may decrease the oral bioavailability of ethinyl estradiol, decreasing the effectiveness of hormonal contraceptives.

Drug-Food Interactions

There were no clinically relevant differences in absorption when ZYDELIG was administered either with food or in a fasting state. Idelalisib can be administered without regard to food.

Drug-Herb Interactions

Coadministration with strong CYP3A inducers such as St. John's Wort should be avoided.

Drug-Laboratory Interactions

Interactions of ZYDELIG with laboratory tests have not been studied.

DOSAGE AND ADMINISTRATION

Dosing Considerations

ZYDELIG can be taken with or without food.

Continue treatment until disease progression or unacceptable toxicity.

Recommended Dose and Dosage Adjustment

Chronic Lymphocytic Leukemia

The recommended dose of ZYDELIG is 150 mg administered orally twice daily in combination with rituximab (8 cycles of rituximab, first cycle at 375 mg/m², subsequent cycles at 500 mg/m²). See the RITUXAN Product Monograph for information on administration and dose adjustment. See Table 5 for ZYDELIG dose modifications.

Follicular Lymphoma

The recommended dose of ZYDELIG is 150 mg administered orally twice daily. See Table 5 for ZYDELIG dose modifications.

Table 5 Dose Modifications for Toxicities Due to ZYDELIG

Event	Grade 1-2	Grade 3	Grade 4
Gastrointestinal			
Diarrhea/Colitis	For Grade 1, provide anti-diarrheal (e.g., loperamide) and maintain ZYDELIG dose. For Grade 2, withhold ZYDELIG and monitor at least weekly until resolved to Grade ≤1.	Withhold ZYDELIG. Consider addition of anti-inflammatory agent (e.g., sulfasalazine, budesonide). Monitor at least weekly until resolved to Grade ≤1, then may resume ZYDELIG at 100 mg BID.	
Hematologic			
Neutropenia	Maintain ZYDELIG dose.	Maintain ZYDELIG dose. Monitor ANC at least weekly.	Interrupt ZYDELIG. Monitor ANC at least weekly until ANC ≥ 0.5 G/L, then may resume ZYDELIG at 100 mg BID.
Hepatic			
ALT/AST Elevation	Maintain ZYDELIG dose. Monitor at least weekly until ALT/AST are ≤1 x ULN.	Withhold ZYDELIG. Monitor at least weekly until ALT/AST are ≤1 x ULN, then may resume ZYDELIG at 100 mg BID.	
Infections			
Evidence of CMV infection or viremia	Discontinue ZYDELIG in patients with evidence of CMV infection or viremia (positive PCR or antigen test).		

Event	Grade 1-2	Grade 3	Grade 4
Evidence of PML	Withhold ZYDELIG at the first sign or symptom suggestive of PML. Discontinue ZYDELIG permanently in patients with confirmed PML.		
Evidence of PCP/PJP Infection	Discontinue ZYDELIG in patients with evidence of PCP/PJP infection.		
Respiratory			
Pneumonitis	Interrupt ZYDELIG and evaluate for respiratory symptoms: <ul style="list-style-type: none"> • If pneumonitis with non-infectious etiology or association with ZYDELIG treatment is suspected, discontinue treatment. • If pneumonitis with infectious etiology is established, monitor until resolved, then may resume ZYDELIG at 100 mg BID. • Permanently discontinue ZYDELIG in patients with evidence of organizing pneumonia. 		
Skin			
Rash	For Grade 1, maintain ZYDELIG dose. For Grade 2, withhold ZYDELIG until Grade ≤ 1 .	Withhold ZYDELIG.	Monitor at least weekly until resolved to Grade ≤ 1 , then may resume ZYDELIG at 100 mg BID.
Abbreviations: ALT, alanine aminotransferase; AST, aspartate aminotransferase; BID, twice daily; CMV, cytomegalovirus; PCP/PJP, <i>Pneumocystis carinii/jirovecii</i> pneumonia; PCR, polymerase chain reaction; ULN, upper limit of normal			

Special Patient Populations

Geriatrics (≥ 65 years of age)

No specific dose adjustment is required for elderly patients (aged ≥ 65 years) (see **ACTION AND CLINICAL PHARMACOLOGY**).

Pediatrics (<18 years of age)

ZYDELIG is not indicated for use in pediatric patients < 18 years of age.

Renal Impairment

No dose adjustment is required for patients with mild, moderate, or severe renal impairment (see **ACTION AND CLINICAL PHARMACOLOGY**).

Hepatic Impairment

No dose adjustment is necessary when initiating treatment with ZYDELIG in patients with mild or moderate hepatic impairment (see **ACTION AND CLINICAL PHARMACOLOGY**). Patients with baseline ALT or AST values greater than 2.5 x ULN or bilirubin values greater than 1.5 x ULN were excluded from pivotal trials. There are insufficient data to make dose recommendations for patients with severe hepatic impairment.

Missed Dose

If a patient misses a dose of ZYDELIG within 6 hours of the time it is usually taken, the patient should take ZYDELIG as soon as possible, and then take the next dose of ZYDELIG at the regularly scheduled time.

If a patient misses a dose of ZYDELIG by more than 6 hours, the patient should not take the missed dose, but resume the usual dosing schedule.

OVERDOSAGE

If overdose occurs, the patient must be monitored for evidence of toxicity. Treatment of overdose with ZYDELIG consists of general supportive measures including monitoring of vital signs as well as observation of the clinical status of the patient.

For management of a suspected drug overdose, please contact your regional Poison Control Centre.
--

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Idelalisib selectively inhibits PI3K δ kinase, which is hyperactive in B-cell malignancies and is central to multiple signaling pathways that drive proliferation, survival, homing, and retention of malignant cells in lymphoid tissues and bone marrow. Idelalisib is a selective inhibitor of adenosine-5'-triphosphate (ATP) binding to the catalytic domain of PI3K δ , resulting in inhibition of the phosphorylation of the key lipid second messenger phosphatidylinositol (PIP) and prevention of Akt phosphorylation.

Idelalisib induces apoptosis and inhibits proliferation in cell lines derived from malignant B-cells and in primary tumor cells. Idelalisib inhibits homing and retention of malignant B-cells in the tumor microenvironment including lymphoid tissues and the bone marrow.

Pharmacodynamics

Effects on Electrocardiogram:

The effect of idelalisib at therapeutic (150 mg) and suprathreshold (400 mg) doses on the QTc interval was evaluated in a placebo- and positive-controlled (moxifloxacin 400 mg) crossover study in 40 healthy subjects. No significant changes in the baseline-corrected QTc based on Fridericia's correction method (QTcF) (i.e., ≥ 10 ms) were observed.

Lymphocytosis:

Upon initiation of ZYDELIG, a temporary increase in lymphocyte counts (i.e., $\geq 50\%$ increase from baseline and above absolute lymphocyte count of 5,000/mcL) has been observed. The onset of isolated lymphocytosis typically occurs during the first two weeks of ZYDELIG therapy. This observed lymphocytosis is a pharmacodynamic effect and should not be considered progressive disease in the absence of other clinical findings.

Pharmacokinetics

Absorption:

Following oral administration of a single 400 mg dose of idelalisib, peak plasma concentrations were observed 2 to 4 hours post-dose under fed conditions and 0.5 to 1.5 hours under fasted conditions.

The C_{max} and AUC of idelalisib increased in a less than dose proportional manner.

Distribution:

Idelalisib is 93% to 94% bound to human plasma proteins at concentrations observed clinically. The mean blood-to-plasma ratio was approximately 0.5.

Metabolism and Elimination:

The metabolism of idelalisib is primarily via aldehyde oxidase, and to a lesser extent via CYP3A and UGT1A4. The primary and only circulating metabolite, GS-563117, is inactive against PI3K δ , but is a strong inhibitor of CYP3A.

The terminal elimination half-life of idelalisib is 8.2 hours following idelalisib 150 mg twice daily oral administration. Following a single 150 mg oral dose of [14 C]-labeled idelalisib, approximately 78% and 15% was excreted in feces and urine, respectively.

Drug-drug Interactions

In vitro, idelalisib inhibited the transport activities of BCRP, OATP1B1 and OATP1B3. A clinical drug interaction study was conducted with rosuvastatin, a sensitive substrate of these transporters. Co-administration of ZYDELIG at 150 mg BID with a single dose of rosuvastatin resulted in comparable rosuvastatin plasma exposures (AUC 90% CI: 87, 121) as observed without ZYDELIG, demonstrating that *in vitro* inhibition of BCRP or OATP1B1/1B3 is not clinically relevant.

Idelalisib inhibited the transport activity of P-gp *in vitro*. In a clinical drug interaction study, digoxin plasma exposures (AUC and C_{max}) were comparable when a single dose of digoxin was administered alone or in combination with ZYDELIG at 150 mg BID, suggesting no clinically relevant inhibition of P-gp or impact on digoxin pharmacokinetics (AUC 90% CI: 98, 111) by ZYDELIG. A risk for P-gp inhibition in the gastrointestinal tract, which could result in increased exposure of sensitive substrates for intestinal P-gp such as dabigatran etexilate, cannot be excluded.

Idelalisib is not an inhibitor of the metabolizing enzymes CYP1A2, CYP2B6, CYP2C, CYP2D6, CYP3A, or UGT1A1, or of the transporters P-gp, BCRP, OATP1B1, OATP1B3, OAT1, OAT3, or OCT2.

GS-563117 is an irreversible (mechanism-based) inhibitor of CYP3A ($K_i = 0.18 \mu\text{M}$, $k_{inact} = 0.033 \text{ min}^{-1}$). GS-563117 is not an inhibitor of the metabolizing enzymes CYP1A2, CYP2B6, CYP2C, CYP2D6 or UGT1A1, or of the transporters P-gp, BCRP, OATP1B1, OATP1B3, OAT1, OAT3, or OCT2.

Special Populations and Conditions

Pediatrics:

The pharmacokinetics of idelalisib has not been studied in pediatric patients.

Geriatrics:

Population pharmacokinetic analyses indicated that age had no clinically relevant effect on the exposures of idelalisib or its primary metabolite GS-563117, including geriatric (65 years of age and older) compared to younger subjects.

Race:

Population pharmacokinetic analyses indicated that race had no clinically relevant effect on the exposures of idelalisib or its primary metabolite GS-563117.

Gender:

Population pharmacokinetic analyses indicated that gender had no clinically relevant effect on the exposures of idelalisib or its primary metabolite GS-563117.

Hepatic Impairment:

A study of pharmacokinetics and safety of idelalisib was performed in healthy volunteers and volunteers with moderate (Child-Pugh Class B) or severe (Child-Pugh Class C) hepatic impairment. Following a single 150 mg dose, no clinically relevant changes in plasma

exposure to idelalisib or its primary metabolite, GS-563117, were observed compared to healthy control volunteers.

Renal Impairment:

A study of pharmacokinetics and safety of idelalisib was performed in healthy volunteers and volunteers with severe renal impairment (estimated creatinine clearance 15 to 29 mL per min). Following a single 150 mg dose, no clinically relevant changes in exposures to idelalisib or its primary metabolite, GS-563117, were observed in subjects with severe renal impairment compared to healthy volunteers.

STORAGE AND STABILITY

Store below 30 °C (86 °F).

- Dispense only in original container.
- Do not use if seal over bottle opening is broken or missing.

SPECIAL HANDLING INSTRUCTIONS

There are no special handling instructions.

DOSAGE FORMS, COMPOSITION AND PACKAGING

ZYDELIG is available as tablets. Each tablet contains 100 or 150 mg of idelalisib. The tablets also include the following inactive ingredients: microcrystalline cellulose, hydroxypropyl cellulose, croscarmellose sodium, sodium starch glycolate and magnesium stearate.

The 150 mg tablets are coated with a material containing red iron oxide, polyethylene glycol, talc, polyvinyl alcohol, and titanium dioxide. ZYDELIG 150 mg tablets are pink, oval-shaped film-coated tablet debossed with “GSI” on one side and the number “150” on the other side. Each bottle contains 60 film-coated tablets and a polyester coil and is closed with a child-resistant closure.

The 100 mg tablets are coated with a material containing Sunset Yellow FCF Aluminum Lake (FD&C Yellow #6), polyethylene glycol, talc, polyvinyl alcohol, and titanium dioxide. ZYDELIG 100 mg tablets are orange, oval-shaped and film-coated, debossed with “GSI” on one side and the number “100” on the other side. Each bottle contains 60 film-coated tablets and a polyester coil and is closed with a child-resistant closure.

PART II. SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

ZYDELIG is the brand name for idelalisib, an isoform-selective, small-molecule inhibitor of phosphatidylinositol 3-kinase p110 δ (PI3K δ).

ZYDELIG tablets are for oral administration. Each tablet contains 150 mg of idelalisib or 100 mg of idelalisib. The tablets also include the following inactive ingredients: microcrystalline cellulose, hydroxypropyl cellulose, croscarmellose sodium, sodium starch glycolate, and magnesium stearate. The 150 mg tablets are film-coated with a material containing polyvinyl alcohol, polyethylene glycol, titanium dioxide, talc, and iron oxide red. The 100 mg tablets are film-coated with a material containing polyvinyl alcohol, polyethylene glycol, titanium dioxide, talc and sunset yellow FCF.

Drug Substance


Common Name: idelalisib

Chemical Name: 5-fluoro-3-phenyl-2-[(1S)-1-(9H-purin-6-ylamino)propyl]quinazolin-4(3H)-one

Empirical Formula: C₂₂H₁₈FN₇O

Molecular Weight: 415.42

Structural Formula:



Physicochemical Properties:

Description: white to off-white solid

Solubility: <0.1 mg/mL at pH 5-7 to over 1 mg/mL at pH 2 under ambient conditions

CLINICAL TRIALS

Chronic Lymphocytic Leukemia

The pivotal CLL study was a randomized, double-blind, placebo-controlled study in 220 patients with relapsed CLL who required treatment but were not considered suitable for cytotoxic chemotherapy based on one of the following criteria: Cumulative Illness Rating Score (CIRS)* >6; estimated CrCl <60 mL/min; Grade ≥3 neutropenia or Grade ≥3 thrombocytopenia resulting from myelotoxic effects of prior therapy with cytotoxic agents. Subjects were randomized 1:1 to receive 8 cycles of rituximab (first cycle at 375 mg/m², subsequent cycles at 500 mg/m²) in combination with either an oral placebo twice daily or with ZYDELIG 150 mg taken twice daily until disease progression or unacceptable toxicity.

The median age was 71 (range 47, 92) with 78.2% of subjects over 65, 65.5% were male, 90.0% were white, 64.5% had a Rai stage of III or IV, and 55.9% had Binet Stage C. Subjects had a median CIRS score of 8; 81 (36.8%) had cardiac, 114 (51.8%) had respiratory, 87 (39.5%) had renal, and 93 (42.3%) had endocrine/metabolic comorbidities. Two hundred and six subjects (93.6%) had 3 or more organs with comorbidities and 82 (37.3%) had severe (score of 3 or higher in any system) comorbidities. The median number of prior therapies was 3.0 (range 1-12). Nearly all (95.9%) subjects had received prior anti-CD20 monoclonal antibodies. The most common (>15%) prior regimens were: bendamustine + rituximab (98 subjects, 44.5%), fludarabine + cyclophosphamide + rituximab (75 subjects, 34.1%), single-agent rituximab (67 subjects, 30.5%), fludarabine + rituximab (37 subjects, 16.8%), and chlorambucil (36 subjects, 16.4%). Most subjects had adverse cytogenetic prognostic factors: 43.2% had a 17p deletion and/or *TP53* mutation, and 83.6% had an unmutated *IGHV*.

The primary endpoint was progression free survival (PFS), defined as the interval from randomization to the earlier of the first documentation of definitive progressive disease (PD) or death from any cause; definitive disease progression was based on standard criteria other than lymphocytosis alone. Other efficacy outcomes included the overall response rate (ORR) and overall survival (OS). The primary analyses of PFS and ORR and were based on evaluation by an independent review committee (IRC).

* The CIRS (Refs) is an index to assess the medical burden of comorbid conditions with total scores ranging from 0 to a theoretical maximum of 56 with higher scores indicating more (or more severe) co-morbidities. Individual comorbidities are assessed in 14 organ systems; each comorbidity is rated with a score from 1-4. If there is >1 comorbidity in an organ system, only the comorbidity with the highest score counts towards the total CIRS score. Conditions that are commonly found to be related to CLL (ie, cytopenia or hypogammaglobulinemia) should not be counted.

The trial was stopped for efficacy following the first pre-specified interim analysis. Results of the second interim analysis continued to show a statistically significant improvement for ZYDELIG + rituximab over placebo + rituximab for the primary endpoint of PFS (HR: 0.18, $p < 0.0001$; see Table 6). This improvement was consistently demonstrated across all pre-specified subgroups (see Figure 2). Nineteen patients died through the cut-off date for the interim analysis; 6 in the ZYDELIG + rituximab group and 13 in the placebo + rituximab group. A statistically significant improvement in ORR was also observed. The Kaplan-Meier plot for PFS is provided in Figure 1.

Table 6. Efficacy Results from Study 312-0116

	ZYDELIG + R n=110	Placebo + R n=110
PFS Median (months) (95% CI)	NR (10.7, NR)	5.5 (3.8, 7.1)
Hazard ratio (95% CI)	0.18 (0.10, 0.32)	
P-value	< 0.0001 [†]	
ORR*, n(%)	82 (74.5%)	16 (14.5%)
Odds ratio (95% CI)	17.28 (8.66, 34.46)	
P-value	< 0.0001 [†]	

R: rituximab; PFS: progression-free survival; NR: not reached

* ORR defined as the proportion of subjects who achieved a CR or PR based on criteria described by Hallek (2008) as modified by Cheson.

† Actual p-values: for PFS, $p = 6 \times 10^{-11}$; for ORR, $p = 6.3 \times 10^{-19}$

Figure 1. Kaplan-Meier Plot of IRC Assessed PFS





Figure 2. Forest Plot of PFS Across Subgroups Favors ZYDELIG + Rituximab



Follicular Lymphoma

The safety and efficacy of ZYDELIG was assessed in a single-arm, multicenter clinical trial that included 72 patients with follicular lymphoma who failed to respond or who had relapsed within 6 months of both rituximab therapy and an alkylating agent (separately or in combination). Subjects received ZYDELIG 150 mg taken orally twice daily until evidence of disease progression or unacceptable toxicity. Tumor response was assessed according to the revised International Working Group response criteria for malignant lymphoma. The primary endpoint was IRC-assessed overall response rate (ORR) summarized in Table 7.

The median age was 62 years (range 33 to 84), 54.2% were male, and 88.9% were Caucasian. At enrollment, 91.7% of patients had a baseline ECOG performance status of 0 or 1. The median time since diagnosis was 4.7 years and the median number of prior treatments was 4 (range 2 to 12). The most common prior combination regimens were R-CHOP (48.6%), BR (48.6%), R (40.3%) and R-CVP (27.8%). At baseline, 33% of patients had extranodal involvement and 26% had bone marrow involvement. Twenty-one patients (29.2%) had disease classified as Grade 1, 39 (54.2%) had disease classified as Grade 2, and 12 (16.7%) had disease classified as Grade 3a.


Table 7. Overall Response Rate (ORR) in Subjects with Follicular Lymphoma

	Number of Subjects (%) N=72
Overall Response (ORR)*	40 (55.6)
95% CI	(43.4, 67.3)
Response Category*	
CR	12 (16.7)
PR	28 (38.9)

* Response as determined by an independent review committee (IRC) where ORR = complete response (CR) + partial response (PR)

The median DOR was 11.8 months (see Figure 3). Of the subjects who did not respond, 23 (31.9%) had stable disease, 8 (11.1%) had progressive disease, and 1 (1.4%) was not evaluable.

Figure 3. Kaplan-Meier Plot of IRC Assessed Duration of Response



DETAILED PHARMACOLOGY

Mechanism of Action

Idelalisib selectively inhibits PI3K δ kinase, which is hyperactive in B-cell malignancies and is central to multiple signaling pathways that drive proliferation, survival, homing, and retention of malignant cells in lymphoid tissues and bone marrow. Idelalisib is a selective inhibitor of adenosine-5'-triphosphate (ATP) binding to the catalytic domain of PI3K δ , resulting in inhibition of the phosphorylation of the key lipid second messenger phosphatidylinositol (PIP) and prevention of Akt phosphorylation.

Idelalisib induces apoptosis and inhibits proliferation in cell lines derived from malignant B-cells and in primary tumor cells. Idelalisib inhibits homing and retention of malignant B-cells in the tumor microenvironment including lymphoid tissues and the bone marrow.

SAFETY PHARMACOLOGY

TOXICOLOGY

Carcinogenesis

The carcinogenicity potential of idelalisib was evaluated in a 26-week transgenic RasH2 mouse study and a 2-year rat study. Idelalisib was not carcinogenic at exposures up to 1.4 and 7.9-fold (by AUC) in male and female mice, respectively compared to the exposure in patients with hematologic malignancies administered the recommended dose of 150 mg twice daily. A dose-dependent increase in pancreatic islet cell tumors was observed at low incidence in male rats at exposures up to 0.4-fold by AUC compared to the human exposure at the recommended dose; a similar finding was not observed in female rats at a 0.62-fold exposure margin.

Mutagenesis

Idelalisib did not induce mutations in the microbial mutagenesis (Ames) assay, was not clastogenic in the *in vitro* chromosome aberration assay using human peripheral blood lymphocytes. Idelalisib was genotoxic in male rats in the *in vivo* micronucleus assay at the highest dose of 2000 mg/kg.

Chronic Toxicity

In rats and dogs, the liver (hepatocellular necrosis), lymphoid tissues (lymphoid depletion) and the male reproductive system (hypospermatogenesis) were identified as target organs/tissues of toxicity in studies conducted up to 26 weeks in rats and 39 weeks in dogs. In dogs, serum transaminase elevations correlated with hepatic necrosis in studies up to 4 weeks; changes appeared to be transient and were not observed following chronic

administration. Changes to lymphoid tissues (spleen, thymus, Peyer's patches) resulted from exaggerated pharmacologic activity of idelalisib.

In rats and dogs, changes in target organs/tissues were either completely or partially reversible following the non-dosing recovery period. In rats, these changes occurred at dose levels where plasma exposure to idelalisib was higher than that observed clinically in patients taking 150 mg idelalisib BID (≥ 2.5 times based on AUC). In dogs, these changes were noted at plasma levels below that observed in humans (≥ 0.25 times based on AUC).

Idelalisib's primary metabolite, GS-563117, has not been qualified in nonclinical repeat dose toxicology studies. Exposure to GS-563117 in rats and dogs in chronic toxicity studies were below that observed clinically in patients taking 150 mg BID.

Safety Pharmacology

There were no idelalisib-related acute effects on CNS function in rats at doses up to 150 mg/kg (approximately 9.2 times human exposure based on C_{max}).

Idelalisib did not produce any acute adverse effects on cardiovascular or respiratory function in dogs up to idelalisib doses of 20 mg/kg (approximately 4.1 times human exposure based on C_{max}). There was no treatment-related prolongation of QT_c interval observed at any dose level.

Reproductive Toxicology

In pregnant rats treated with idelalisib at 25, 75, or 150 mg/kg/day for 12 days (gestation day 6 to 17), postimplantation loss, lower mean fetal weights and skeletal development variations were observed at 75 and 150 mg/kg/day were observed (≥ 12 times human exposure based on AUC).

Idelalisib was embryotoxic and teratogenic at dose levels inducing maternal toxicity in rats. Maternal toxicity was demonstrated by dose-dependent decreases in the body weight gains of the dams. Dose-dependent developmental findings included higher incidence of postimplantation loss, decrease in viable fetuses, and decreased mean fetal body weights. Dose-dependent external malformations included those of skeletal origin consistent with vertebral agenesis and short tails. Additional external malformations were single instances of hydrocephaly and microphthalmia, occurring in separate fetuses from different litters.

Fertility

The male reproductive system was a target organ of idelalisib toxicity in both rats and dogs. Idelalisib may impair fertility in humans. In male rats treated with idelalisib at 25, 50, or 100 mg/kg/day for 10 weeks, decreases in epididymides and testes weight were observed

but with no adverse effects on mating or fertility parameters, and no degeneration or loss in spermatogenesis (≤ 7.8 times human exposure based on AUC).

Phototoxicity

Results in the *in vitro* 3T3 NRU phototoxicity assay were inconclusive for idelalisib due to cytotoxicity in the assay. GS-563117 may induce phototoxicity in the presence of UVA exposure.

REFERENCES

1. Furman R, et al. Idelalisib and Rituximab in Relapsed Chronic Lymphocytic Leukemia. *N Engl J Med* 2014; DOI: 10.1056/NEJMoa1315226.
2. Gopal A, et. al. PI3K δ Inhibition by Idelalisib in Patients with Relapsed Indolent Lymphoma. *N Engl J Med* 2014; DOI: 10.1056/NEJMoa1314583.
3. Extermann M. Measuring Comorbidity in Older Cancer Patients. *Eur J of Cancer* 2000, 36, 453-471.
4. Parmelee P, et al. Validation of the Cumulative Illness Rating Scale in a Geriatric Residential Population. *J Amer Geriat Soc* 1995, 43:130-137.

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

**PrZYDELIG®
idelalisib tablets**

Read this carefully before you start taking ZYDELIG and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about ZYDELIG.

Serious Warnings and Precautions

Take ZYDELIG only under the care of a qualified doctor who knows how to use drugs to treat cancer.

During treatment and for 2 to 6 months after you stop taking ZYDELIG:

- take antibiotics to prevent a type of pneumonia called Pneumocystis.
- continue to be monitored for a viral infection called cytomegalovirus.

Hepatotoxicity (liver damage) is a serious side effect.

Serious side effects that can lead to death include:

- Serious infections.
- Severe diarrhea and colitis (digestive disease).
- Pneumonitis (severe lung problems).
- Stevens-Johnson Syndrome and Toxic Epidermal Necrolysis (severe skin reactions).

What is ZYDELIG used for?

ZYDELIG is used in adults to treat the following types of cancers:

- **Chronic Lymphocytic Leukemia (CLL):** ZYDELIG is used in combination with rituximab to treat a blood and bone marrow cancer. It is called chronic lymphocytic leukemia (CLL). ZYDELIG is used when you have been previously treated for your cancer.
- **Follicular Lymphoma (FL):** ZYDELIG is used alone to treat a blood cancer. It is called follicular lymphoma (FL). ZYDELIG is used when your cancer did not respond to at least two previous treatments, including rituximab and an alkylating agent.

ZYDELIG is NOT for use in children under the age of 18 years.

How does ZYDELIG work?

ZYDELIG blocks a specific protein in the body that helps cancer cells live and grow. This protein is called "PI3 Kinase". By blocking this protein, ZYDELIG may help kill and reduce the number of cancer cells and slow the spread of the cancer in your body.

What are the ingredients in ZYDELIG?

Medicinal ingredients: idelalisib.

Non-medicinal ingredients: croscarmellose sodium, hydroxypropyl cellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, and sodium starch glycolate, talc, and titanium dioxide.

100 mg orange tablets also have: FD&C Yellow #6/Sunset Yellow, and FCF Aluminum Lake.

150 mg pink tablets also have: red iron oxide

ZYDELIG comes in the following dosage forms:

Tablets: 100 mg, 150 mg

Do not use ZYDELIG if:

- if you are allergic to idelalisib or any of the other ingredients in this product.

ZYDELIG should not be prescribed by your healthcare professional as the first medicine to treat CLL or as the first or second medicine to treat FL.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take ZYDELIG. Talk about any health conditions or problems you may have, including if you:

- have an infection from any cause such as bacteria, fungus or virus. Serious infections can happen while taking ZYDELIG. Your doctor will prescribe a medicine and check your blood to reduce the risk of getting certain infections.
- have liver damage, hepatitis or liver disease. Before and while taking ZYDELIG, you will have regular blood tests to check your liver. Your doctor may decide to stop treatment, or briefly stop treatment to let your liver get better, before starting the treatment again at a lower dose.
- have severe diarrhea. Diarrhea is common while taking ZYDELIG and can sometimes be severe. Tell your doctor right away if the number of bowel movements you have in a day increases to four or more times. Ask your doctor about medicines you can take to treat your diarrhea.
- have breathing or severe lung problems. Your doctor may do tests to check your lungs if you have breathing problems while taking ZYDELIG.
- have severe skin reactions.

Pregnancy, birth control and breastfeeding:

- ZYDELIG should not be used during pregnancy. While on ZYDELIG do NOT get pregnant. ZYDELIG can harm your unborn baby.
- Use a reliable method of birth control while taking ZYDELIG and for 1 month after you stop taking ZYDELIG. Talk to your doctor about the birth control you may use.
- ZYDELIG may make the birth control pill or oral contraceptive not work as well. You must also add a barrier method to be sure you do not get pregnant while taking ZYDELIG and for 1 month after your last treatment.
- Tell your doctor right away if you become pregnant while taking ZYDELIG.
- Do not breastfeed while taking ZYDELIG. If you are currently breastfeeding, talk to your doctor before starting treatment. It is not known if ZYDELIG passes into human milk.

Other warnings you should know about:

Progressive Multifocal Leukoencephalopathy (PML):

- Cases of PML have been reported following ZYDELIG use. PML is a rare brain infection that can be fatal. Tell your doctor right away if you notice or someone notices in you: progressive weakness on one side of the body, clumsiness of limbs, disturbance of vision, changes in thinking, memory and orientation, confusion, personality changes. Your doctor may request further testing if PML is suspected. Your doctor should stop treatment if PML is confirmed.

Serious Allergic Reactions:

- Serious allergic reactions can occur while taking ZYDELIG. Get medical help right away if you experience symptoms of an allergic reaction. These include rash, hives, swelling of the face, lips, tongue or throat or difficulty swallowing or breathing. Be sure to tell your doctor if this has happened to you. Your doctor may decide to stop treatment.

Sensitivity to Sunlight:

- You may become sensitive to the sun while taking ZYDELIG. You should stay out of the sun or wear sunscreen.

Driving and Using Machines:

- ZYDELIG is not likely to affect you being able to drive or use any tools or machines. Use caution until you know how ZYDELIG affects you.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with ZYDELIG:

- alfentanil, fentanyl (used for pain relief).
- amiodarone, disopyramide, lidocaine, quinidine (used to treat heart problems).
- amlodipine, diltiazem, felodipine, nifedipine (used to treat high blood pressure and heart problems).
- atorvastatin, lovastatin, simvastatin (used to lower cholesterol).
- buspirone, clorazepate, diazepam, flurazepam, zolpidem (used to treat nervous system disorders).
- carbamazepine, phenytoin, S-mephenytoin (used to prevent seizures).
- cyclosporine, sirolimus, tacrolimus (used to control your body's immune response after a transplant).
- dihydroergotamine, ergotamine (used to treat migraines or headaches).
- ketoconazole (used to treat fungal infections).
- midazolam, triazolam (used to help you sleep and/or relieve anxiety).
- pimozide (used to treat illnesses affecting the way you think, feel or behave).
- quetiapine (used to treat schizophrenia, bipolar disorder and major depressive disorder).
- quinidine (used to treat irregular heartbeats).
- raloxifene (used to treat or prevent bone loss).
- repaglinide (used to lower blood sugar).
- rifampicin (used to prevent and treat tuberculosis and other infections).
- St. John's wort (*Hypericum perforatum*, an herbal product used for anxiety or depression).
- sildenafil, tadalafil (used to treat impotence and pulmonary hypertension, a lung disease that

- makes breathing difficult).
- oral or implanted hormonal contraceptives (used to prevent pregnancy).
- warfarin (used to thin the blood).

How to take ZYDELIG:

- Take ZYDELIG every day. Follow the instructions from your doctor, exactly as it says on the label. Set up a dosing schedule and follow it carefully. Talk to your healthcare professional if you have any questions.
- Take this medicine with or without food.
- Your doctor will tell you the dose of rituximab that is right for you.
- Do not change or stop your treatment without first talking with your doctor. Take ZYDELIG every day until your doctor tells you to stop. This happens if your cancer no longer responds to the drug or you have unacceptable side effects.

It is important to see your doctor regularly while taking ZYDELIG. Only take medicine that has been prescribed to you. Do not give ZYDELIG to others.

Recommended Adult Dose: one 150 mg tablet twice a day.

Your doctor may reduce your dose of ZYDELIG to 100 mg twice a day if you experience side effects.

Overdose:

If you think you have taken too much ZYDELIG, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

It is important to take ZYDELIG each day.

- If you miss a dose of ZYDELIG and it is **less than 6 hours** from the time you usually take ZYDELIG, take the dose right away.
- If **more than 6 hours** has passed from the time you usually take ZYDELIG, wait and take the next dose at your usual time. Do NOT take a double dose (two doses close together).

What are possible side effects from using ZYDELIG?

These are not all the possible side effects you may feel when taking ZYDELIG. If your side effect is not listed here, contact your doctor. Side effects may occur at any time, even months or a year after starting or stopping ZYDELIG.

Side effects may include:

- diarrhea, nausea, vomiting.
- heartburn, stomach pain, decreased appetite.
- sores/ulcers in the mouth and/or on the lips.
- headache.
- fatigue, feeling tired.
- trouble sleeping.
- stuffy nose, colds, sinus infections, bronchitis, cough.

- rash.
- chills.
- night sweats.
- muscle pain.
- joint pain.

ZYDELIG can cause abnormal blood test results, including, but not limited to, high and low blood sugar. Your doctor will decide when to perform blood tests and will read the results. You will need a blood test at least every 2 weeks for the first 6 months on ZYDELIG.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
VERY COMMON			
Hepatotoxicity (liver damage): yellowing of your skin or eyes, dark urine, pain in your abdomen, nausea, vomiting, loss of appetite		✓	
Fever	✓		
Severe diarrhea and colitis (digestive disease): increased number of bowel movements, watery or bloody stool, stomach pain and/or cramps		✓	
Neutropenia (low numbers of white blood cells): fever, chills or sweating, or any signs of infection		✓	
Serious infections: fever, sweats, or chills, cough or flu-like symptoms like rapid breathing, shortness of breath, blood in your phlegm, muscle aches, pain in your stomach			✓
COMMON			
Pneumonitis or pneumonia (severe lung problems caused by a fungus called <i>Pneumocystis</i> or a virus called cytomegalovirus): new or worsening cough, shortness of breath, difficult or painful breathing, wheezing, pain in chest when breathing, fever		✓	

Severe skin reactions (including Stevens-Johnson syndrome, toxic epidermal necrolysis, and drug reaction with eosinophilia and systemic symptoms [DRESS]): skin rash that spreads quickly, redness, fever, enlarged lymph nodes, blistering and peeling of the skin and/or inside of the lips, eyes, mouth, nasal passages or genitals			✓
Peripheral edema: swelling, especially of the ankles and feet		✓	
Urinary tract infection: strong and frequent urge to urinate, cloudy, bloody or strong smelling urine, pain or burning sensation when urinating		✓	
UNCOMMON			
Serious allergic reactions: rash, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			✓
Progressive multifocal leukoencephalopathy (PML) (a rare brain infection): progressive weakness on one side of the body, clumsiness of limbs, disturbance of vision, changes in thinking, memory and orientation, confusion, personality changes			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Store below 30 °C (86 °F).
- Do not use ZYDELIG if the seal over the bottle opening is broken or missing.
- Keep out of reach and sight of children.

If you want more information about ZYDELIG:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website (www.gilead.ca), or by calling 1-800-207-4267.

This leaflet was prepared by Gilead Sciences Canada, Inc.

Revised: 18 August 2020

Gilead Sciences, Inc.

Foster City, CA 94404
USA

Gilead Sciences Canada, Inc.

Mississauga, ON L5N 2W3

ZYDELIG® is a trademark of Gilead Sciences, Inc. or its related companies.

All other marks referenced herein are property of their respective owners.

© 2020 Gilead Sciences, Inc. All rights reserved.



e165440-GS-006